

Geometric Interpretation of the Function $g(c, \mu)$ of Neutron Transport Theory

Perry A. Newman

NASA Langley Research Center
Hampton, Virginia 23365

and

Robert L. Bowden

Department of Physics, Virginia Polytechnic Institute
Blacksburg, Virginia 24061

Received January 23, 1970

Revised April 3, 1970

In their classic monograph on neutron transport theory, Case et al.¹ introduced the function $g(c, \mu)$,

$$g(c, \mu) = [(1 - c\mu \tanh^{-1} \mu)^2 + (\pi c\mu/2)^2]^{-1}, \quad (1)$$

and gave many of its properties. It is now well known² that this function is simply related to the normalization of the singular eigenfunctions introduced later by Case.³ An analogous function is readily defined² in time-dependent monoenergetic neutron-transport problems in plane geometry with isotropic scattering; however it also contains a complex transform variable. Bowden and Williams⁴ have shown that in such time-dependent problems the discrete eigenfunctions are not always present in the normal-mode expansion of the Laplace transform of the solution; that is, there exists a curve, C_s , in the s/c -plane (see Fig. 1) such that if s/c lies inside the curve ($s \in S_i$) discrete terms are present, whereas for s/c outside the curve ($s \in S_e$) there are no discrete terms. The curve C_s is given by⁴

$$C_s = \left\{ s/c = \alpha' + i\beta' \mid \alpha' = \frac{2\beta'}{\pi} \tanh^{-1} \frac{2\beta'}{\pi} \right\}. \quad (2)$$

We want to point out in this note that the function

$$\frac{1}{\alpha^2} g\left(\frac{1}{\alpha}, \mu\right) = \frac{c^2}{\Omega^+(\mu, s)\Omega^-(\mu, s)} \Big|_{\beta=0}, \quad 0 \leq \mu \leq 1, \quad (3)$$

where $s/c = \alpha + i\beta$ and $\Omega^\pm(\mu, s)$ are the limiting values of the dispersion function

$$\Omega(z, s) = s - cz \tanh^{-1}(1/z) \quad (4)$$

on its branch cut in the z -plane, is the inverse of the square of the distance in the s/c -plane from the point $(\alpha, 0)$ to the point $[\alpha'(\mu), \beta'(\mu)]$ which lies on the curve C_s .

The functions $|\Omega^\pm(\mu, s)/c|^2$ are easily found from Eq. (4) to be

$$\left| \frac{\Omega^\pm(\mu, s)}{c} \right|^2 = (\alpha - \mu \tanh^{-1} \mu)^2 + (\beta \pm \pi\mu/2)^2, \quad (5)$$

while the parametric form of Eq. (2) is

$$|\beta'(\mu)| = \pi\mu/2, \quad \alpha'(\mu) = \mu \tanh^{-1} \mu, \quad 0 \leq \mu \leq 1. \quad (6)$$

Thus $|\Omega^\pm(\mu, s)/c|^2$ are the squares of the distances in the s/c -plane from the point (α, β) to the points $[\alpha'(\mu), \pm\beta'(\mu)]$ which lie on the curve C_s , in the lower and upper half-planes, respectively. When $\beta = 0$, $\Omega^+(\mu, s)$ and $\Omega^-(\mu, s)$ are complex conjugates so that Eq. (5) reduces to Eq. (3) and the stated result follows.




Fig. 1. Location of the curve C_s in the complex s/c -plane, $s/c = \alpha + i\beta$. Several values of the parameter μ are given in parentheses for C_s in the upper half-plane. The function $\nu_0(s)$ is the solution of $\Omega(\nu_0, s) = 0$ for which $\text{Re}(\nu_0) > 0$ when $\text{Re}(s/c) > 1$. Some values of $\nu_0(s)$ are plotted in the complex s -plane in Fig. 1 of Ref. 5 (Fig. 7.1 of Ref. 2).

Case et al. show, for example, that $g(1/\alpha, \mu)|_{\max}$ occurs at $\mu = 0$ for $\alpha < \pi^2/8$, whereas for $\alpha > \pi^2/8$ it occurs for μ between 0 and 1. For α very large, they show that $g_{\max} \rightarrow 4\alpha^2/\pi^2$. The present geometric interpretation is seen to be consistent with these characteristics. The radius of curvature of the curve C_s given by Eq. (2) is $\pi^2/8$ at $(\alpha', \beta') = (0, 0)$ as indicated by the dashed circle on Fig. 1. For α' very large, $\beta' \rightarrow \pi/2$ so that the minimum squared distance from $(\alpha, 0)$ to (α', β') approaches $\pi^2/4$, in agreement with Eq. (3) and $g_{\max} \rightarrow 4\alpha^2/\pi^2$.

¹K. M. CASE, F. deHOFFMAN, and G. PLACZEK, *Introduction to the Theory of Neutron Diffusion*, U.S. Government Printing Office, Washington, D.C. (1953).

²K. M. CASE and P. F. ZWEIFEL, *Linear Transport Theory*, Addison-Wesley Pub. Co., Reading, Massachusetts (1967).

³K. M. CASE, *Ann. Phys.*, **9**, 1 (1960).

⁴R. L. BOWDEN and C. D. WILLIAMS, *J. Math. Phys.*, **5**, 1527 (1964).

⁵I. KUŠČER and P. F. ZWEIFEL, *J. Math. Phys.*, **6**, 1125 (1965).